

Package ‘climetrics’

July 19, 2022

Type Package

Title Climate Change Metrics

Version 1.0-5

Date 2022-07-18

Depends methods, sp ($\geq 1.2-0$), R ($\geq 3.5.0$), rts

Imports raster, terra, zoo, yalImpute, xts

Suggests R.rsp

VignetteBuilder R.rsp

Description A framework that facilitates spatio-temporal analysis of climate dynamics through exploring and measuring different dimensions of climate change in space and time.

License GPL (≥ 3)

Encoding UTF-8

URL <http://r-gis.net>

BugReports <https://github.com/shirintaheri/climetrics/issues/>

NeedsCompilation no

Author Shirin Taheri [cre, aut] (<https://orcid.org/0000-0002-0303-3145>),
Babak Naimi [aut] (<https://orcid.org/0000-0001-5431-2729>),
Miguel Araujo [aut]

Maintainer Shirin Taheri <taheri.shi@gmail.com>

Repository CRAN

Date/Publication 2022-07-19 15:00:05 UTC

R topics documented:

aaClimate	2
apply.months	4
ccm	6
daClimate	9
dVelocity	12

gVelocity	13
kgc	14
localExtreme	16
novelClimate	18
sed	19
temporalTrend	21
velocity	22

Index	24
--------------	-----------

aaClimate	<i>Changes in the Area of Analogous Climate</i>
-----------	---

Description

To quantify the change in area of analogous climates (classes), the change in area occupied by a given class between time 1 (t1) and time 2 (t2) periods is quantified.

Usage

```
aaClimate(precip,tmin, tmax, tmean,t1,t2)
aaClimateC(c1,c2)
```

Arguments

precip	A time series of precipitation as a Raster or Raster Time Series object
tmin	A time series of minimum temperature as a Raster or Raster Time Series object
tmax	A time series of maximum temperature as a Raster or Raster Time Series object
tmean	A time series of mean temperature as a Raster or Raster Time Series object; if not provided, it will be calculated from tmin and tmax
t1	a character or a numeric vector, specifying the index of raster layers for time 1
t2	a character or a numeric vector, specifying the index of raster layers for time 2
c1	A single layer Raster layer contains Climate classes (e.g., Koggen Geiger climate classification) for time 1
c2	A single layer Raster layer contains Climate classes (e.g., Koggen Geiger climate classification) for time 2

Details

For a given cell with a given climate class, the change in area of analogous climates represented the ratio (in percentage) of the difference between time 2 and time 1 area of that class to the time 1 area of the same class. Positive values indicated gains in area, negative values indicated losses, and null values reflect no change.

The aaClimate first uses apply.months function to generate the monthly mean of climate parameter (results a Raster object with 12 layers correspond to 12 months) over each time period (t1 and t2); then uses the kgc function to generate the Koggen Geiger climate classification for each time period. Then the changes in the area of each class is calculated between time 1 and time 2. If the climate classification (regions) are available for time 1 and time 2, then the aaClimateC can be used instead.

Value

A single Raster layer (RasterLayer or SpatRaster depending on the input)

Author(s)

Shirin Taheri; Babak Naimi

<taheri.shi@gmail.com>; <naimi.b@gmail.com>

Examples

```
#-----
filePath <- system.file("external/", package="climetrics") # path to the dataset folder

# read the climate variables using the terra package (you can use the raster package as well):

pr <- rast(paste0(filePath, '/precip.tif'))
tmin <- rast(paste0(filePath, '/tmin.tif'))
tmax <- rast(paste0(filePath, '/tmax.tif'))
tmean <- rast(paste0(filePath, '/tmean.tif'))

pr # has 360 layers corresponds to months of the years 1991-2020

n <- readRDS(paste0(filePath, '/dates.rds')) # read corresponding dates

class(n)
length(n)

head(n) # Dates corresponds to the layers in climate variables (pr, tmin, tmax, tmean)

#####

# use rts function in the rts package to make a raster time series:

pr.t <- rts(pr,n)
tmin.t <- rts(tmin,n)
tmax.t <- rts(tmax,n)
tmean.t <- rts(tmean,n)
#-----

pr.t # see the summary report of the raster time series object

#####
# test of the metric:
#-----
#-----
# t1 (time1) = '1991/1990' takes all layers correspond to years between 1991-01-01 to 2000-12-31
# t2 (time2) = '2010/2020' takes all layers correspond to years between 2010-01-01 to 2020-12-31

aa <- aaClimate(precip=pr.t, tmin=tmin.t, tmax=tmax.t, tmean=tmean.t, t1='1991/2000', t2='2010/2020')
```

```

plot(aa,main="Changes in the area of Analogous Climates")

#####
# Alternatively, if the climate in two times are presented as climate classes
# (e.g., Koggen Geiger climate classification), changes in distance to analogous
# climate can be quantified using the daClimateC function:

# Here, we first generate Koggen Geiger climate classification for time 1 and time 2, separately,
# we need monthly average of climate variables (over 12 months):

# take average of climate variables over each of 12 months:

p12.1 <- apply.months(pr.t[['1991/2000']], 'mean')
p12.2 <- apply.months(pr.t[['2010/2020']], 'mean')

p12.1

plot(p12.1)
#--
tmin12.1 <- apply.months(tmin.t[['1991/2000']], 'mean')
tmin12.2 <- apply.months(tmin.t[['2010/2020']], 'mean')
#--
tmax12.1 <- apply.months(tmax.t[['1991/2000']], 'mean')
tmax12.2 <- apply.months(tmax.t[['2010/2020']], 'mean')
#--
tmean12.1 <- apply.months(tmean.t[['1991/2000']], 'mean')
tmean12.2 <- apply.months(tmean.t[['2010/2020']], 'mean')
#--

##----- now, the kgc function can be used to generate the climate clasification map:
k1 <- kgc(p=p12.1,tmin = tmin12.1,tmax=tmax12.1, tmean = tmean12.1)
k2 <- kgc(p=p12.2,tmin = tmin12.2,tmax=tmax12.2, tmean = tmean12.2)

plot(k1, main= "Koggen Geiger climate classification - 1995")
plot(k2, main= "Koggen Geiger climate classification - 2015")

aa <- aaClimateC(k1,k2)

plot(aa)

```

Description

Apply a user specified function over raster time series for each month separately.

Usage

```
apply.months(x, FUN, ...)
```

Arguments

x	A Raster or Raster Time Series object
FUN	A function to apply on rasters of each month over different years; The defaults if mean
...	additional argument: if x is a Raster or SpatRaster object, corresponding dates should be specified in the dates argument

Details

The output contains 12 raster layers corresponding to the 12 months (Jan-Dec.). For each of the available 12 months (from January to December), the corresponding rasters are selected and the FUN is applied. For example if the FUN is mean, the first layer in the output would be the mean of the rasters in January over different years, and the last later (12th layer) would be the mean of the rasters in December over different years.

Value

A Raster object (RasterStackBrick or SpatRaster depending on the input) contains 12 layers correspond to 12 months (From Jan to Dec)

Author(s)

Shirin Taheri; Babak Naimi
<taheri.shi@gmail.com>; <naimi.b@gmail.com>

Examples

```
filePath <- system.file("external/", package="climetrics") # path to the dataset folder

pr <- rast(paste0(filePath, '/precip.tif'))

n <- readRDS(paste0(filePath, '/dates.rds')) # read corresponding dates

head(n) # Dates corresponds to the layers in climate variables (pr, tmin, tmax, tmean)

#####

# use rts function in the rts package to make a raster time series:

pr.t <- rts(pr, n)
```

```
#####

p12 <- apply.months(pr.t, 'mean')

p12

plot(p12)
```

ccm

Climate Change Metrics

Description

A function to quantify a single or multiple climate change metrics between two time periods given the climatic raster time series variables.

Usage

```
ccm(x, ..., stat, t1, t2, extreme, longlat, ny, names, verbose=TRUE)
```

Arguments

x	a Raster object or a Raster Time Series of a climate variable
...	additional climate variables can be included as a Raster object or a Raster Time Series
stat	a character specifying the metric used to generate climate change metrics, can be one or multiple metrics including 'sed', 'novelClimate', 'localExtreme', 'aaClimate', 'daClimate', 'velocity', 'gVelocity', 'dVelocity' (or their abbreviations)
t1	a character or a numeric vector, specifying the index of raster layers for time 1
t2	a character or a numeric vector, specifying the index of raster layers for time 2
extreme	the percentile used for stat='lce', local climate extreme event change
longlat	TRUE or FALSE, specifies whether the input data is geographic or projected
ny	integer; specifies the number of years between time 1 and time2, if the input is Raster time series, it will be identified by the function
names	character with a length equal to the number of input climate variables; specifies the names of the input climate variables; for some metrics the names (precipitation, tmin, tmax, tmean) is required
verbose	logical (default=TRUE), specifies whether the function print messages as a text output on the screen

Details

multiple climate change metrics are available to quantify multiple dimensions of climate change between two times.

The metrics include:

- sed: Standardized Local Anomalies
- localExtreme (lce): Changes in probability of local extremes
- aaClimate (aac): Changes in areas of analogous climates
- novelClimate (nc): Novel climates
- daClimate (dac): Changes in distance to analogous climates
- velocity (ve): Climate change velocity (based on the method represented in the paper Hamnan et al. (2015))
- dVelocity (dVe): Distance-based climate change velocity (based on the method used in Garcia et al. (2014))
- gVelocity (gVe): Gradient-based climate change velocity (based on the method represented in Burrow et al. (2011))

References:

- Burrows MT, Schoeman DS, Buckley LB, Moore P, Poloczanska ES, Brander KM, Brown C, Bruno JF, Duarte CM, Halpern BS, Holding J. (2011) The pace of shifting climate in marine and terrestrial ecosystems. *Science*. Nov 4;334(6056):652-5.
- Garcia, R.A., Cabeza, M., Rahbek, C. and Araújo, M.B. (2014). Multiple dimensions of climate change and their implications for biodiversity. *Science*, 344(6183).
- Hamann, A., Roberts, D.R., Barber, Q.E., Carroll, C. and Nielsen, S.E. (2015). Velocity of climate change algorithms for guiding conservation and management. *Global Change Biology*, 21(2), pp.997-1004.

Value

A Raster object (RasterLayer or RasterStackBrick or SpatRaster depending on the input) with single or multiple layers correspond to the selected metrics (stat)

Author(s)

Shirin Taheri; Babak Naimi

<taheri.shi@gmail.com>; <naimi.b@gmail.com>

<http://r-gis.net>

Examples

```
#-----
filePath <- system.file("external/", package="climetrics") # path to the dataset folder

# read the climate variables using the terra package (you can use the raster package as well):
```

```

pr <- rast(paste0(filePath, '/precip.tif'))
tmin <- rast(paste0(filePath, '/tmin.tif'))
tmax <- rast(paste0(filePath, '/tmax.tif'))
tmean <- rast(paste0(filePath, '/tmean.tif'))

pr # has 360 layers corresponds to months of the years 1991-2020

# n <- names(pr)
# n <- substring(n,2,11)
# head(n)
# n <- as.Date(n,format = '%Y.%m.%d')

n <- readRDS(paste0(filePath, '/dates.rds')) # read corresponding dates

class(n)
length(n)

head(n) # Dates corresponds to the layers in climate variables (pr, tmin, tmax, tmean)

#####

# use rts function in the rts package to make a raster time series:

pr.t <- rts(pr,n)
tmin.t <- rts(tmin,n)
tmax.t <- rts(tmax,n)
tmean.t <- rts(tmean,n)
#-----

pr.t # see the summary report of the raster time series object

#####
# test of the metric:
#-----
#-----
# t1 (time1) = '1991/1990' takes all layers correspond to years between 1991-01-01 to 2000-12-31
# t2 (time2) = '2010/2020' takes all layers correspond to years between 2010-01-01 to 2020-12-31

sed <- ccm(pr.t,tmin.t,tmean.t,tmax.t,t1='1991/2000',t2='2010/2020',stat='sed')

plot(sed, main='Standardized Local Anomalies')

s2 <- ccm(pr.t,tmin.t,tmean.t,tmax.t,t1='1991/2000',t2='2010/2020',stat=c('nc','sed'))

plot(s2)

ve <- ccm(pr.t,tmin.t,tmean.t,tmax.t,t1='1991/2000',t2='2010/2020',stat=c('gve','dve'))

```



```

plot(ve)

# following, the extreme argument is needed for localExtreme (lce) metric:
s3 <- ccm(pr.t,tmin.t,t1='1991/2000',t2='2010/2020',stat=c('localExtreme','sed')
          ,extreme = c(0.05,0.95))

plot(s3)

```

daClimate

Changes in distance to Analogous Climates

Description

Quantifies the changes in distances to analogous climates between two time periods.

Usage

```

daClimate(precip,tmin, tmax, tmean,t1,t2)
daClimateC(c1,c2)

```

Arguments

precip	A time series of precipitation as a Raster or Raster Time Series object
tmin	A time series of minimum temperature as a Raster or Raster Time Series object
tmax	A time series of maximum temperature as a Raster or Raster Time Series object
tmean	A time series of mean temperature as a Raster or Raster Time Series object; if not provided, it will be calculated from tmin and tmax
t1	a character or a numeric vector, specifying the index of raster layers for time 1
t2	a character or a numeric vector, specifying the index of raster layers for time 2
c1	A single layer Raster layer contains Climate classes (e.g., Koggen Geiger climate classification) for time 1
c2	A single layer Raster layer contains Climate classes (e.g., Koggen Geiger climate classification) for time 2

Details

For a given cell, the function quantifies the distances to all cells with analogous climates in the time 1 period, i.e., belonging to the same climate class of the given cell. It also quantifies the distances to all cells that experience analogous climates in the time 2. Then, for each cell, it calculates the median of the great-circle distances (in km) below the 10th percentile of the distribution of all values, for both time 1 and time 2 periods, and maps the change over time. Negative values indicated a temporal decrease in distance, whereas positive values indicated an increase.

The daClimate first uses `apply.months` function to generate the monthly mean of climate parameter (results a Raster object with 12 layers correspond to 12 months) over each time period (t1 and t2); then uses the `kgc` function to generate the Koggen Geiger climate classification for each time period. Then the changes in the area of each class is calculated between time 1 and time 2. If the climate classification (regions) are available for time 1 and time 2, then the `daClimateC` can be used instead.

Value

A single Raster layer (RasterLayer or SpatRaster depending on the input)

Author(s)

Shirin Taheri; Babak Naimi

<taheri.shi@gmail.com>; <naimi.b@gmail.com>

Examples

```
#-----
filePath <- system.file("external/", package="climetrics") # path to the dataset folder

# read the climate variables using the terra package (you can use the raster package as well):

pr <- rast(paste0(filePath, '/precip.tif'))
tmin <- rast(paste0(filePath, '/tmin.tif'))
tmax <- rast(paste0(filePath, '/tmax.tif'))
tmean <- rast(paste0(filePath, '/tmean.tif'))

pr # has 360 layers corresponds to months of the years 1991-2020

n <- readRDS(paste0(filePath, '/dates.rds')) # read corresponding dates

class(n)
length(n)

head(n) # Dates corresponds to the layers in climate variables (pr, tmin, tmax, tmean)

#####

# use rts function in the rts package to make a raster time series:

pr.t <- rts(pr, n)
```

```

tmin.t <- rts(tmin,n)
tmax.t <- rts(tmax,n)
tmean.t <- rts(tmean,n)
#-----

pr.t # see the summary report of the raster time series object

#####
# test of the metric:
#-----
#-----
# t1 (time1) = '1991/1990' takes all layers correspond to years between 1991-01-01 to 2000-12-31
# t2 (time2) = '2010/2020' takes all layers correspond to years between 2010-01-01 to 2020-12-31

da <- daClimate(precip=pr.t,tmin=tmin.t,tmax=tmax.t,tmean=tmean.t,t1='1991/2000',t2='2010/2020')

plot(da,main="Changes in distance to Analogous Climates")

#####
# Alternatively, if the climate in two times are presented as climate classes
# (e.g., Koggen Geiger climate classification), changes in distance to analogous
# climate can be quantified using the daClimateC function:

# Here, we first generate Koggen Geiger climate classification for time 1 and time 2, separately,
# we need monthly average of climate variables (over 12 months):

# take average of climate variables over each of 12 months:

p12.1 <- apply.months(pr.t[['1991/2000']], 'mean')
p12.2 <- apply.months(pr.t[['2010/2020']], 'mean')

p12.1

plot(p12.1)
#--
tmin12.1 <- apply.months(tmin.t[['1991/2000']], 'mean')
tmin12.2 <- apply.months(tmin.t[['2010/2020']], 'mean')
#--
tmax12.1 <- apply.months(tmax.t[['1991/2000']], 'mean')
tmax12.2 <- apply.months(tmax.t[['2010/2020']], 'mean')
#--
tmean12.1 <- apply.months(tmean.t[['1991/2000']], 'mean')
tmean12.2 <- apply.months(tmean.t[['2010/2020']], 'mean')
#--

##----- now, the kgc function can be used to generate the climate classification map:
k1 <- kgc(p=p12.1,tmin = tmin12.1,tmax=tmax12.1, tmean = tmean12.1)
k2 <- kgc(p=p12.2,tmin = tmin12.2,tmax=tmax12.2, tmean = tmean12.2)

plot(k1, main= "Koggen Geiger climate classification - 1995")
plot(k2, main= "Koggen Geiger climate classification - 2015")

```

```
da <- daClimateC(k1,k2)

plot(da)
```

dVelocity

Distant-based Velocity of Climate Change

Description

The method is developed based on the approach used in Garcia et al. (2014) Gradient-based velocity is also implemented in the package and is available through the gVelocity function.

Garcia, R.A., Cabeza, M., Rahbek, C. and Araújo, M.B. (2014). Multiple dimensions of climate change and their implications for biodiversity. *Science*, 344(6183).

Usage

```
dVelocity(x,...,t1, t2, ny)
```

Arguments

x	The first input climate variable that can be a RasterStack or RasterBrick, or a raster time series
...	additional input climate variables in case of using multiple climate variables that can be entered as v1, v2, ...
t1	a character or a numeric vector, specifying the index of raster layers for time 1
t2	a character or a numeric vector, specifying the index of raster layers for time 2
ny	integer; specifies the number of years between time 1 and time2, if the input is Raster time series, it is not needed as it will be identified by the function

Value

A single Raster layer (RasterLayer or SpatRaster depending on the input)

Author(s)

Shirin Taheri; Babak Naimi

<taheri.shi@gmail.com>; <naimi.b@gmail.com>

Examples

```

filePath <- system.file("external/", package="climetrics") # path to the dataset folder

pr <- rast(paste0(filePath, '/precip.tif'))
tmean <- rast(paste0(filePath, '/tmean.tif'))

n <- readRDS(paste0(filePath, '/dates.rds')) # corresponding dates

head(n) # Dates corresponds to the layers in climate variables

#####

# use rts function in the rts package to make a raster time series:

pr.t <- rts(pr,n)
tmean.t <- rts(tmean,n)
#####

dv <- dVelocity(pr.t,tmean.t,t1='1991/2000',t2='2010/2020')

plot(dv)

```

gVelocity

*Gradient-based Velocity of Climate Change***Description**

The method is developed based on Burrow et al. (2011) which is a gradient-based velocity of climate change. Climate velocity can inform conservation planning and action in a warming world.

Distance-based velocity is also implemented in the package and is available through the ccm function.

Burrows MT, Schoeman DS, Buckley LB, Moore P, Poloczanska ES, Brander KM, Brown C, Bruno JF, Duarte CM, Halpern BS, Holding J. The pace of shifting climate in marine and terrestrial ecosystems. *Science*. 2011 Nov 4;334(6056):652-5..

Usage

```
gVelocity(x, ...)
```

Arguments

x	a Raster object or a Raster Time Series of climate variable
...	additional arguments; not implemented

Value

A single Raster layer (RasterLayer or SpatRaster depending on the input)

Author(s)

Shirin Taheri; Babak Naimi
<taheri.shi@gmail.com>; <naimi.b@gmail.com>

Examples

```
filePath <- system.file("external/", package="climetrics") # path to the dataset folder
# read the climate variables using the terra package (you can use the raster package as well):
pr <- rast(paste0(filePath, '/precip.tif'))
gv <- gVelocity(pr)
plot(gv, main='Gradient-based Velocity for Precipitation')
```

kgc

Koggen Geiger climate classification

Description

Given the monthly mean (12 months) of Precipitation, minimum, maximum, and mean temperature datasets, this function identify climate zones following Koggen Geiger climate classification rules.

Usage

```
kgc(p, tmin, tmax, tmean)
```

Arguments

p	A Raster object contains 12 months precipitation
tmin	A Raster object contains 12 months minimum temperature
tmax	A Raster object contains 12 months maximum temperature
tmean	A Raster object contains 12 months mean temperature

Details

If the input object is not a 12 month Raster object but a time series, the function first calls the `apply.months` to get the 12 month raster object.

Value

A single Raster layer (RasterLayer or SpatRaster depending on the input)

Author(s)

Shirin Taheri; Babak Naimi

<taheri.shi@gmail.com>; <naimi.b@gmail.com>

Examples

```

filePath <- system.file("external/", package="climetrics") # path to the dataset folder

# read the climate variables using the terra package (you can use the raster package as well):

pr <- rast(paste0(filePath, '/precip.tif'))
tmin <- rast(paste0(filePath, '/tmin.tif'))
tmax <- rast(paste0(filePath, '/tmax.tif'))
tmean <- rast(paste0(filePath, '/tmean.tif'))

pr # has 360 layers corresponds to months of the years 1991-2020

n <- readRDS(paste0(filePath, '/dates.rds')) # read corresponding dates

class(n)
length(n)

head(n) # Dates corresponds to the layers in climate variables (pr, tmin, tmax, tmean)

#####

# use rts function in the rts package to make a raster time series:

pr.t <- rts(pr,n)
tmin.t <- rts(tmin,n)
tmax.t <- rts(tmax,n)
tmean.t <- rts(tmean,n)
#-----

pr.t # see the summary report of the raster time series object

#####

# To generate Koggen Geiger climate classification, we need monthly average of
# climate variables (over 12 months):

# you can use apply.months function to take the average of climate variables over
# each month that generates 12 layers corresponds to 12 months:

p12 <- apply.months(pr.t[['1991/2000']], 'mean')

```

```

p12

# plot(p12)
#--
tmin12 <- apply.months(tmin.t[['1991/2000']], 'mean')

tmax12 <- apply.months(tmax.t[['1991/2000']], 'mean')

tmean12 <- apply.months(tmean.t[['1991/2000']], 'mean')

#--

##----- now, the kgc function can be used to generate the climate clasification map:
k1 <- kgc(p=p12,tmin = tmin12,tmax=tmax12, tmean = tmean12)

k1

plot(k1)

```

localExtreme

Changes in probability of local extremes

Description

Probability of changes in local extremes is calculated at each cell based on either one or two climate variables (e.g., precipitation and temperature). For each climate variable, the extreme value (percentile of the extreme in the variable distribution) should be specified by user in the extreme parameter. For instance, if temperature and precipitation are specified as the first and second inputs (x_1 and x_2), $\text{extreme}=\text{c}(0.95,0.05)$ may be used that considers 95th and 5th percentiles of the distribution of the temperature and precipitation time series, respectively over the first time period, and then it calculates the probability that the time1 percentile will be exceeded in time2. If two climate variables are used (e.g., temperature and precipitation), for each cell, the two probabilities are summed from which the product of the two probabilities are subtracted to obtain a measure of the probability of occurrence of either of the two extreme events (to avoid counting probabilities twice). The probability of time_1 extreme climates in time_2, is then subtracted from the probability of time_1 extreme climates to obtain the changes in the probability of extreme climate variables. Positive values indicated increased probability in time 2, whereas negative values indicated a decrease.

Usage

```
localExtreme(x1,x2,t1,t2,extreme)
```


Arguments

x1	Time Series of the first climate variable as a Raster or Raster Time Series Object
x2	Time Series of the second climate variable as a Raster or Raster Time Series Object; x1 and x2 can be lists of time series of climate variables in time 1 and time 2, respectively
t1	a character or a numeric vector, specifying the index of raster layers for time 1
t2	a character or a numeric vector, specifying the index of raster layers for time 2
extreme	a numeric vector with a length of one or two (depends on whether one or two climate variables are used as input), specifying the percentile of extreme value in the input (first value corresponds to x1 and second corresponds to x2 climate variables)

Value

A single Raster layer (RasterLayer or SpatRaster depending on the input)

Author(s)

Shirin Taheri; Babak Naimi

<taheri.shi@gmail.com>; <naimi.b@gmail.com>

Examples

```

filePath <- system.file("external/", package="climetrics") # path to the dataset folder

# read the climate variables using the terra package (you can use the raster package as well):

pr <- rast(paste0(filePath, '/precip.tif'))
tmax <- rast(paste0(filePath, '/tmax.tif'))

pr # has 360 layers corresponds to months of the years 1991-2020

n <- readRDS(paste0(filePath, '/dates.rds')) # read corresponding dates

head(n) # Dates corresponds to the layers in climate variables (pr, tmin, tmax, tmean)

#####

# use rts function in the rts package to make a raster time series:

pr.t <- rts(pr,n)
tmax.t <- rts(tmax,n)

#####
# test of the metric:

```

```
# The extreme argument corresponds to the first and second climate variables
# (i.e., x1 and x2; precipitation and temperature) that specify the percentile of the extreme
# condition in climate variable; here, 0.05 is used for precipitation; and 0.95 for temperature

le <- localExtreme(x1=pr.t,x2=tmax.t,t1='1991/2000',t2='2010/2020', extreme = c(0.05, 0.95))

plot(le, main='Probability of Changes in Local Climate Extreme')
```

novelClimate

Novel Climates

Description

Novel climate is one of the climate change metrics to quantify the dissimilarities between climate parameters between two time periods. This metric uses standard euclidean distance between each cell in Time 2 and all cells in Time 1 and retains the minimum of those distances. The inter-annual standard deviation for each parameter is used for the standardization. The larger the score, the more dissimilar the climate in Time 2 is in relation to the global pool of potential climatic analogues.

Usage

```
novelClimate(x, ..., t1, t2)
```

Arguments

x	a Raster object or a Raster Time Series of a climate variable
...	additional climate variables can be included as a Raster object or a Raster Time Series
t1	a character or a numeric vector, specifying the index of raster layers for time 1; if x is missing, t1 and t2 can be list of climate variables in time 1 and time 2, respectively.
t2	a character or a numeric vector, specifying the index of raster layers for time 2; if x is missing, t1 and t2 can be list of climate variables in time 1 and time 2, respectively.

Value

A single Raster layer (RasterLayer or SpatRaster depending on the input)

Author(s)

Shirin Taheri; Babak Naimi
 <taheri.shi@gmail.com>; <naimi.b@gmail.com>

Examples

```

filePath <- system.file("external/", package="climetrics") # path to the dataset folder

# read the climate variables using the terra package (you can use the raster package as well):

pr <- rast(paste0(filePath, '/precip.tif'))
tmean <- rast(paste0(filePath, '/tmean.tif'))

pr # has 360 layers corresponds to months of the years 1991-2020

n <- readRDS(paste0(filePath, '/dates.rds')) # read corresponding dates

head(n) # Dates corresponds to the layers in climate variables (pr, tmin, tmax, tmean)

#####

# use rts function in the rts package to make a raster time series:

pr.t <- rts(pr,n)
tmean.t <- rts(tmean,n)

#####
# test of the metric:

n1 <- novelClimate(pr.t,tmean.t,t1='1991/2000',t2='2010/2020')

plot(n1, main='Novel Climate')

#####
# if the input object is SpatRaster (or RasterStack or RasterBrick) object:
# t1 and t2 would be the numbers specifying which layers correspond to time1 and time2:

n2 <- novelClimate(pr,tmean,t1=1:120,t2=229:360)

plot(n2)

```

Description

Standardized Local Anomalies is one of the climate change metrics to quantify the change between two time periods.

Usage

```
sed(x, ..., t1, t2)
```

Arguments

x	a Raster object or a Raster Time Series of a climate variable
...	additional climate variables can be included as a Raster object or a Raster Time Series
t1	a character or a numeric vector, specifying the index of raster layers for time 1
t2	a character or a numeric vector, specifying the index of raster layers for time 2

Value

A single Raster layer (RasterLayer or SpatRaster depending on the input)

Author(s)

Shirin Taheri; Babak Naimi
<taheri.shi@gmail.com>; <naimi.b@gmail.com>

Examples

```
filePath <- system.file("external/", package="climetrics") # path to the dataset folder
# read the climate variables using the terra package (you can use the raster package as well):
pr <- rast(paste0(filePath, '/precip.tif'))
tmin <- rast(paste0(filePath, '/tmin.tif'))
tmax <- rast(paste0(filePath, '/tmax.tif'))

n <- readRDS(paste0(filePath, '/dates.rds')) # read corresponding dates
head(n) # Dates corresponds to the layers in climate variables (pr, tmin, tmax, tmean)
#####

# use rts function in the rts package to make a raster time series:

pr.t <- rts(pr, n)
tmin.t <- rts(tmin, n)
tmax.t <- rts(tmax, n)

#####
# test of the metric:

s <- sed(pr.t, tmin.t, tmax.t, t1='1991/2000', t2='2010/2020')

plot(s, main='Standardized Local Anomalies')
```

```
#####  
# if the input object is SpatRaster (or RasterStack or RasterBrick) object:  
# t1 and t2 would be the numbers specifying which layers correspond to time1 and time2:  
  
s <- sed(pr, tmin, tmax, t1=1:120, t2=229:360)  
  
plot(s)
```

temporalTrend	<i>Temporal Trend (Slope) of a climate variable</i>
---------------	---

Description

The method calculates the trend of a climate variable changes over time. The function uses a Raster time series as input and returns a Raster object that represents the trend at the pixel level.

Usage

```
temporalTrend(x, ...)
```

Arguments

x	a Raster object or a Raster Time Series of climate variable
...	additional arguments; not implemented

Value

A single Raster layer (RasterLayer or SpatRaster depending on the input)

Author(s)

Shirin Taheri; Babak Naimi
<taheri.shi@gmail.com>; <naimi.b@gmail.com>

Examples

```
filePath <- system.file("external/", package="climetrics") # path to the dataset folder  
  
# read the climate variables using the terra package (you can use the raster package as well):  
pr <- rast(paste0(filePath, '/precip.tif'))  
  
tr <- temporalTrend(pr)
```

```
# plot(tr, main='Trend (slope) of Precipitation time series')
```

 velocity

Velocity of Climate Change

Description

The method is developed based on the method represented in the paper Hamnan et al. (2015). Two additional functions including Distance-based (`dVelocity`) and Gradient-based velocity (`gVelocity`) are also implemented and available in the package.

- Hamann, A., Roberts, D.R., Barber, Q.E., Carroll, C. and Nielsen, S.E. (2015). Velocity of climate change algorithms for guiding conservation and management. *Global Change Biology*, 21(2), pp.997-1004.

Usage

```
velocity(x1,x2,t1, t2,...)
```

Arguments

x1	The first input climate variable that can be a RasterStack or RasterBrick, or a raster time series
x2	The second input climate variable that can be a RasterStack or RasterBrick, or a raster time series
t1	a character or a numeric vector, specifying the index of raster layers for time 1
t2	a character or a numeric vector, specifying the index of raster layers for time 2
...	not implemented

Value

A single Raster layer (RasterLayer or SpatRaster depending on the input)

Author(s)

Shirin Taheri; Babak Naimi

<taheri.shi@gmail.com>; <naimi.b@gmail.com>

Examples

```
filePath <- system.file("external/", package="climetrics") # path to the dataset folder

# read the climate variables using the terra package (you can use the raster package as well):

pr <- rast(paste0(filePath, '/precip.tif'))
tmax <- rast(paste0(filePath, '/tmax.tif'))

pr # has 360 layers corresponds to months of the years 1991-2020

n <- readRDS(paste0(filePath, '/dates.rds')) # read corresponding dates

head(n) # Dates corresponds to the layers in climate variables (pr, tmin, tmax, tmean)

#####

# use rts function in the rts package to make a raster time series:

pr.t <- rts(pr,n)
tmax.t <- rts(tmax,n)

#####

# test of the metric:

# The extreme argument corresponds to the first and second climate variables
# (i.e., x1 and x2; precipitation and temperature) that specify the percentile of the extreme
# condition in climate variable; here, 0.05 is used for precipitation; and 0.95 for temperature

ve <- velocity(x1=pr.t,x2=tmax.t,t1='1991/2000',t2='2010/2020')

# plot(ve, main='Velocity of Climate Change')
```

Index

aaClimate, [2](#)
aaClimate, RasterStackBrick-method
 (aaClimate), [2](#)
aaClimate, RasterStackBrickTS-method
 (aaClimate), [2](#)
aaClimate, SpatRaster-method
 (aaClimate), [2](#)
aaClimate, SpatRasterTS-method
 (aaClimate), [2](#)
aaClimateC (aaClimate), [2](#)
aaClimateC, RasterStackBrick-method
 (aaClimate), [2](#)
aaClimateC, SpatRaster-method
 (aaClimate), [2](#)
apply.months, [4](#)
apply.months, RasterStackBrick-method
 (apply.months), [4](#)
apply.months, RasterStackBrickTS-method
 (apply.months), [4](#)
apply.months, SpatRaster-method
 (apply.months), [4](#)
apply.months, SpatRasterTS-method
 (apply.months), [4](#)

ccm, [6](#)
ccm, RasterStackBrick-method (ccm), [6](#)
ccm, RasterStackBrickTS-method (ccm), [6](#)
ccm, SpatRaster-method (ccm), [6](#)
ccm, SpatRasterTS-method (ccm), [6](#)

daClimate, [9](#)
daClimate, RasterStackBrick-method
 (daClimate), [9](#)
daClimate, RasterStackBrickTS-method
 (daClimate), [9](#)
daClimate, SpatRaster-method
 (daClimate), [9](#)
daClimate, SpatRasterTS-method
 (daClimate), [9](#)
daClimateC (daClimate), [9](#)

daClimateC, RasterStackBrick-method
 (daClimate), [9](#)
daClimateC, SpatRaster-method
 (daClimate), [9](#)
dVelocity, [12](#)
dVelocity, RasterStackBrick-method
 (dVelocity), [12](#)
dVelocity, RasterStackBrickTS-method
 (dVelocity), [12](#)
dVelocity, SpatRaster-method
 (dVelocity), [12](#)
dVelocity, SpatRasterTS-method
 (dVelocity), [12](#)

gVelocity, [13](#)
gVelocity, RasterStackBrick-method
 (gVelocity), [13](#)
gVelocity, RasterStackBrickTS-method
 (gVelocity), [13](#)
gVelocity, SpatRaster-method
 (gVelocity), [13](#)
gVelocity, SpatRasterTS-method
 (gVelocity), [13](#)

kgc, [14](#)
kgc, RasterStackBrick, RasterStackBrick, RasterStackBrick-method
 (kgc), [14](#)
kgc, SpatRaster, SpatRaster, SpatRaster-method
 (kgc), [14](#)

localExtreme, [16](#)
localExtreme, list, list, ANY, ANY-method
 (localExtreme), [16](#)
localExtreme, list, list-method
 (localExtreme), [16](#)
localExtreme, missing, missing, list, list-method
 (localExtreme), [16](#)
localExtreme, RasterStackBrick, ANY, ANY, ANY-method
 (localExtreme), [16](#)

- localExtreme,RasterStackBrick-method
(localExtreme), [16](#)
- localExtreme,RasterStackBrickTS,ANY,ANY,ANY-method
(localExtreme), [16](#)
- localExtreme,RasterStackBrickTS-method
(localExtreme), [16](#)
- localExtreme,SpatRaster,ANY,ANY,ANY-method
(localExtreme), [16](#)
- localExtreme,SpatRaster-method
(localExtreme), [16](#)
- localExtreme,SpatRasterTS,ANY,ANY,ANY-method
(localExtreme), [16](#)
- localExtreme,SpatRasterTS-method
(localExtreme), [16](#)

- novelClimate, [18](#)
- novelClimate,missing-method
(novelClimate), [18](#)
- novelClimate,RasterStackBrick-method
(novelClimate), [18](#)
- novelClimate,RasterStackBrickTS-method
(novelClimate), [18](#)
- novelClimate,SpatRaster-method
(novelClimate), [18](#)
- novelClimate,SpatRasterTS-method
(novelClimate), [18](#)

- sed, [19](#)
- sed,missing-method (sed), [19](#)
- sed,RasterStackBrick-method (sed), [19](#)
- sed,RasterStackBrickTS-method (sed), [19](#)
- sed,SpatRaster-method (sed), [19](#)
- sed,SpatRasterTS-method (sed), [19](#)

- temporalTrend, [21](#)
- temporalTrend,RasterStackBrick-method
(temporalTrend), [21](#)
- temporalTrend,RasterStackBrickTS-method
(temporalTrend), [21](#)
- temporalTrend,SpatRaster-method
(temporalTrend), [21](#)
- temporalTrend,SpatRasterTS-method
(temporalTrend), [21](#)

- velocity, [22](#)
- velocity,RasterStackBrick,RasterStackBrick-method
(velocity), [22](#)
- velocity,RasterStackBrickTS,RasterStackBrickTS-method
(velocity), [22](#)
- velocity,SpatRaster,SpatRaster-method
(velocity), [22](#)
- velocity,SpatRasterTS,SpatRasterTS-method
(velocity), [22](#)